

ESTROGENIC COMPOUNDS  
AS ANTI-MITOTIC AGENTS

Background of the Invention

5 This invention relates to treating disease states characterized by abnormal cell mitosis.

Cell mitosis is a multi-step process that includes cell division and replication (Alberts, B. et al. In *The Cell*, pp. 652-661 (1989); Stryer, E. *Biochemistry* (1988)).

10 Mitosis is characterized by the intracellular movement and segregation of organelles, including mitotic spindles and chromosomes. Organelle movement and segregation are facilitated by the polymerization of the cell protein tubulin. Microtubules are formed from  $\alpha$  and  $\beta$  tubulin  
15 polymerization and the hydrolysis of GTP. Microtubule formation is important for cell mitosis, cell locomotion, and the movement of highly specialized cell structures such as cilia and flagella.

Microtubules are extremely labile structures that  
20 are sensitive to a variety of chemically unrelated anti-mitotic drugs. For example, colchicine and nocadazole are anti-mitotic drugs that bind tubulin and inhibit tubulin polymerization (Stryer, E. *Biochemistry* (1988)). When used alone or in combination with other therapeutic drugs,  
25 colchicine may be used to treat cancer (WO-9303729-A, published March 4, 1993; J03240726-A, published October 28, 1991), alter neuromuscular function, change blood pressure, increase sensitivity to compounds affecting sympathetic neuron function, depress respiration, and relieve gout  
30 (*Physician's Desk Reference*, Vol. 47, p. 1487, (1993)).

Estradiol and estradiol metabolites such as 2-methoxyestradiol have been reported to inhibit cell division (Seegers, J.C. et al. *J. Steroid Biochem.* 32, 797-809 (1989); Lottering, M-L. et al. *Cancer Res.* 52, 5926-5923

1007742-024002

205720" 2472001

(1992); Spicer, L.J. and Hammond, J.M. *Mol. and Cell. Endo.* 64, 119-126 (1989); Rao, P.N. and Engelberg, J. *Exp. Cell Res.* 48, 71-81 (1967)). However, the activity is variable and depends on a number of *in vitro* conditions. For  
5 example, estradiol inhibits cell division and tubulin polymerization in some *in vitro* settings (Spicer, L.J. and Hammond, J.M. *Mol. and Cell. Endo.* 64, 119-126 (1989); Ravindra, R., *J. Indian Sci.* 64(c) (1983)), but not in  
10 others (Lottering, M-L. et al. *Cancer Res.* 52, 5926-5923 (1992); Ravindra, R., *J. Indian Sci.* 64(c) (1983)). Estradiol metabolites such as 2-methoxyestradiol will inhibit cell division in selected *in vitro* settings depending on whether the cell culture additive phenol red is  
15 present and to what extent cells have been exposed to estrogen. (Seegers, J.C. et al. Joint NCI-IST Symposium. Biology and Therapy of Breast Cancer. 9/25-9/27, 1989, Genoa, Italy, Abstract A58).

Numerous diseases are characterized by abnormal cell mitosis. For example, uncontrolled cell mitosis is a  
20 hallmark of cancer. In addition, cell mitosis is important for the normal development of the embryo, formation of the corpus luteum, wound healing, inflammatory and immune responses, angiogenesis and angiogenesis related diseases.

#### Summary of the Invention

25 ~~I have discovered that certain compounds within the scope of the general formulae set forth below in the claims are useful for treating mammalian diseases characterized by undesired cell mitosis. Without wishing to bind myself to any particular theory, such compounds generally inhibit  
30 microtuble formation and tubulin polymerization and/or depolymerization. Compounds within the general formulae having said inhibiting activity are preferred. Preferred compositions may also exhibit a change (increase or~~

decrease) in estrogen receptor binding, improved absorption, transport (e.g. through blood-brain barrier and cellular membranes), biological stability, or decreased toxicity. I have also discovered certain compounds useful in the method, as described by the general formulae of the claims.

A mammalian disease characterized by undesirable cell mitosis, as defined herein, includes but is not limited to excessive or abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying: rheumatoid arthritis, psoriasis, diabetic retinopathy, and other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neovascular glaucoma and Osler Weber syndrome. Other undesired angiogenesis involves normal processes including ovulation and implantation of a blastula. Accordingly, the compositions described above can be used to block ovulation and implantation of a blastula or to block menstruation (induce amenorrhea).

The bond indicated by C...C is absent or, in combination with the C---C bond is the unit HC=CH.

Other features and advantages of the invention will be apparent from the following description of preferred embodiments thereof.

~~Description of the Preferred Embodiments~~  
~~The drawings are first described.~~

Fig. 1 is a graph illustrating the inhibition of tubulin polymerization by 2-methoxyestradiol described by Example 1 below.

Fig. 2 is a graph illustrating the inhibition of colchicine binding to tubulin by 2-methoxyestradiol described by Example 2 below.

Fig. 3 depicts: I. colchicine, 2-methoxyestradiol  
5 and combretastatin A-4, and II. various estradiol derivatives comprising colchicine (a-c) or combretastatin A-4 (d) structural motifs as described below.

Compounds According to the Invention

As described below, compounds that are useful in  
10 accordance with the invention include novel estradiol derivatives that bind tubulin, inhibit microtubule formation or exhibit anti-mitotic properties. Specific compounds according to the invention are described below. Those  
15 skilled in the art will appreciate that the invention extends to other compounds within the formulae given in the claims below, having the described characteristics. These characteristics can be determined for each test compound using the assays detailed below and elsewhere in the literature.

Without wishing to bind myself to specific  
20 mechanisms or theory, it appears that certain compounds that are known to inhibit microtubule formation, bind tubulin and exhibit anti-mitotic properties such as colchicine and combretastatin A-4 share certain structural similarities  
25 with estradiol. Fig. 3 illustrates the molecular formulae of estradiol, colchicine, combretastatin A-4, and improved estradiol derivatives that bind tubulin inhibit microtubule assembly and exhibit anti-mitotic properties. Molecular formulae are drawn and oriented to emphasize structural  
30 similarities between the ring structures of colchicine, combretastatin A-4, estradiol, and certain estradiol derivatives. Estradiol derivatives are made by

incorporating ~~colchicine~~ or combretastatin A-4 structural motifs into the steroidal backbone of estradiol.

Figure 3, part I, depicts the chemical formulae of colchicine, 2-methoxyestradiol and combretastatin A-4.

- 5 Figure 3, part IIa-d, illustrates estradiol derivatives that comprise structural motifs found in colchicine or combretastatin A-4. For example, part II a-c shows estradiol derivatives with an A and/or B ring expanded from six to seven carbons as found in colchicine and part II d depicts an estradiol derivative with a partial B ring as found in combretastatin A-4. Each C ring of an estradiol derivative, including those shown in Figure 3, may be fully saturated as found in 2-methoxyestradiol. R<sub>1-6</sub> represent a subset of the substitution groups found in the claims. Each
- 10 R<sub>1-6</sub> can independently be defined as -R<sub>1</sub>, OR<sub>1</sub>, -OCOR<sub>1</sub>, -SR<sub>1</sub>, -F, -NHR<sub>2</sub>, -Br, -I, or -C≡CH.

*In situ*  
Anti-mitotic Activity In Situ

- Anti-mitotic activity is evaluated *in situ* by testing the ability of an improved estradiol derivative to
- 20 inhibit the proliferation of new blood vessel cells (angiogenesis). A suitable assay is the chick embryo chorioallantoic membrane (CAM) assay described by Crum et al. *Science* 230:1375 (1985). See also, U.S. Patent 5,001,116, hereby incorporated by reference, which describes
- 25 the CAM assay. Briefly, fertilized chick embryos are removed from their shell on day 3 or 4, and a methylcellulose disc containing the drug is implanted on the chorioallantoic membrane. The embryos are examined 48 hours later and, if a clear avascular zone appears around the
- 30 methylcellulose disc, the diameter of that zone is measured. Using this assay, a 100mg disk of the estradiol derivative 2-methoxyestradiol was found to inhibit cell mitosis and the growth of new blood vessels after 48 hours. This result

indicates that the anti-mitotic action of 2-methoxyestradiol can inhibit cell mitosis and angiogenesis.

#### Anti-Mitotic Activity In Vitro

Anti-mitotic activity can be evaluated by testing the ability of an estradiol derivative to inhibit tubulin polymerization and microtubule assembly *in vitro*. Microtubule assembly is followed in a Gilford recording spectrophotometer (model 250 or 2400S) equipped with electronic temperature controllers. A reaction mixture (all concentrations refer to a final reaction volume of 0.25 $\mu$ l) contains 1.0M monosodium glutamate (ph 6.6), 1.0mg/ml (10 $\mu$ M) tubulin, 1.0 mM MgCl<sub>2</sub>, 4% (v/v) dimethylsulfoxide and 20-75 $\mu$ M of a composition to be tested. The 0.24ml reaction mixtures are incubated for 15 min. at 37°C and then chilled on ice. After addition of 10 $\mu$ l 2.5mM GTP, the reaction mixture is transferred to a cuvette at 0°C, and a baseline established. At time zero, the temperature controller of the spectrophotometer is set at 37°C. Microtubule assembly is evaluated by increased turbidity at 350 nm. Alternatively, inhibition of microtubule assembly can be followed by transmission electron microscopy as described in Example 2 below.

#### Indications

The invention can be used to treat any disease characterized by abnormal cell mitosis. Such diseases include, but are not limited to: abnormal stimulation of endothelial cells (e.g., atherosclerosis), solid tumors and tumor metastasis, benign tumors, for example, hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, vascular malfunctions, abnormal wound healing, inflammatory and immune disorders, Bechet's disease, gout or gouty arthritis, abnormal angiogenesis accompanying: rheumatoid arthritis, psoriasis, diabetic retinopathy, and

other ocular angiogenic diseases such as retinopathy of prematurity (retrolental fibroplastic), macular degeneration, corneal graft rejection, neurovascular glaucoma and Oster Webber syndrome.

5 Improved Estradiol Derivative Synthesis

Known compounds that are used in accordance with the invention and precursors to novel compounds according to the invention can be purchased, e.g., from Sigma Chemical Co., St. Louis, Steroloids and Research Plus. Other compounds  
10 according to the invention can be synthesized according to known methods from publicly available precursors.

The chemical synthesis of estradiol has been described (Eder, V. et al., Ber 109, 2948 (1976); Oppolzer, D.A. and Roberts, D.A. *Helv. Chim. Acta.* 63, 1703, (1980)).  
15 Synthetic methods for making seven-membered rings in multi-cyclic compounds are known (Nakamura, T. et al. *Chem. Pharm. Bull.* 10, 281 (1962); Sunagawa, G. et al. *Chem. Pharm. Bull.* 9, 81 (1961); Van Tamelen, E. E. et al. *Tetrahedron* 14, 8-34 (1961); Evans, D. E. et al. *JACS* 103, 5813 (1981)). Those  
20 skilled in the art will appreciate that the chemical synthesis of estradiol can be modified to include 7-membered rings by making appropriate changes to the starting materials, so that ring closure yields seven-membered rings. Estradiol or estradiol derivatives can be modified to  
25 include appropriate chemical side groups according to the invention by known chemical methods (*The Merck Index*, 11th Ed., Merck & Co., Inc., Rahway, NJ USA (1989), pp. 583-584).  
Administration

The compositions described above can be provided as  
30 physiologically acceptable formulations using known techniques, and these formulations can be administered by standard routes. In general, the combinations may be administered by the topical, oral, rectal or parenteral

205720 247400

(e.g., intravenous, subcutaneous or intramuscular) route. In addition, the combinations may be incorporated into biodegradable polymers allowing for sustained release, the polymers being implanted in the vicinity of where delivery is desired, for example, at the site of a tumor. The biodegradable polymers and their use are described in detail in Brem et al., *J. Neurosurg.* 74:441-446 (1991).

The dosage of the composition will depend on the condition being treated, the particular derivative used, and other clinical factors such as weight and condition of the patient and the route of administration of the compound. However, for oral administration to humans, a dosage of 0.01 to 100 mg/kg/day, preferably 0.01-1 mg/kg/day, is generally sufficient.

The formulations include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intraocular, intratracheal, and epidural) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous



liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion and as a bolus, etc.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients.

- 5 Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tables may be made by  
10 molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally coated or scored and may be formulated so as to provide a slow or controlled release of the active ingredient therein.

- 15 Formulations suitable for topical administration in the mouth include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and  
20 mouthwashes comprising the ingredient to be administered in a suitable liquid carrier.

- Formulations suitable for topical administration to the skin may be presented as ointments, creams, gels and pastes comprising the ingredient to be administered in a  
25 pharmaceutical acceptable carrier. A preferred topical delivery system is a transdermal patch containing the ingredient to be administered.

- Formulations for rectal administration may be presented as a suppository with a suitable base comprising,  
30 for example, cocoa butter or a salicylate.

Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to

10077442-021502  
500 microns which is administered in the manner in which  
snuff is taken, i.e., by rapid inhalation through the nasal  
passage from a container of the powder held close up to the  
nose. Suitable formulations, wherein the carrier is a  
5 liquid, for administration, as for example, a nasal spray or  
as nasal drops, include aqueous or oily solutions of the  
active ingredient.

Formulations suitable for vaginal administration may  
be presented as pessaries, tampons, creams, gels, pastes,  
10 foams or spray formulations containing in addition to the  
active ingredient such as carriers as are known in the art  
to be appropriate.

Formulations suitable for parenteral administration  
include aqueous and non-aqueous sterile injection solutions  
15 which may contain anti-oxidants, buffers, bacteriostats and  
solutes which render the formulation isotonic with the blood  
of the intended recipient; and aqueous and non-aqueous  
sterile suspensions which may include suspending agents and  
thickening agents. The formulations may be presented in  
20 unit-dose or multi-dose containers, for example, sealed  
ampules and vials, and may be stored in a freeze-dried  
(lyophilized) conditions requiring only the addition of the  
sterile liquid carrier, for example, water for injections,  
immediately prior to use. Extemporaneous injection  
25 solutions and suspensions may be prepared from sterile  
powders, granules and tables of the kind previously  
described.

Preferred unit dosage formulations are those  
containing a daily dose or unit, daily sub-dose, as herein  
30 above recited, or an appropriate fraction thereof, of the  
administered ingredient.

*para 17*  
It should be understood that in addition to the  
ingredients, particularly mentioned above, the formulations

of this invention may include other agents convention in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include flavoring agents.

5 Example 1:

Figure 1 illustrates the inhibition of tubulin polymerization by 2-methoxyestradiol.

10 A. Each reaction mixture (all concentrations refer to the final reaction volume of 0.25 ml) contained 1.0 M monosodium glutamate (pH 6.6), 1.0 mg/ml (10  $\mu$ M) tubulin, 1.0 mM  $MgCl_2$ , 4% (v/v) dimethylsulfoxide, and either 0 (curve 1), 20  $\mu$ M (curve 2), 40  $\mu$ M (curve 3), or 75  $\mu$ M (curve 4) 2-methoxyestradiol. The 0.24 ml reaction mixtures were incubated for 15 min at 37°C and chilled on ice. After  
15 addition of 10  $\mu$ l of 2.5 mM GTP the reaction mixtures were transferred to cuvettes held at 0°C, and baselines were established. At time zero the temperature controller was set at 37°C. At the times indicated by the vertical dashed lines the temperature controller was set at the indicated  
20 temperatures.

25 B. Each reaction mixture contained 0.8 M monosodium glutamate (pH 6.6), 1.2 mg/ml (12  $\mu$ M) tubulin, 4% (v/v) dimethylsulfoxide, and either 0 (curve 1), 1.0  $\mu$ M (curve 2), 2.0  $\mu$ M (curve 3), 3.0  $\mu$ M (curve 4), or 4.0  $\mu$ M (curve 5) 2-methoxyestradiol. The 0.24 ml reaction mixtures were incubated for 15 min at 26°C and chilled on ice. After addition of 10  $\mu$ l of 10 mM GTP the reaction mixtures were transferred to cuvettes held at 0°C, and baselines were established. At time zero the temperature controller was  
30 set at 26°C. At the time indicated by vertical dashed line the temperature controller was set at 0°C.

Example 2:

Transmission electron microscopy (TEM) can show differences between the morphology of polymerized tubulin formed in the absence or presence of 2-methoxyestradiol.

5 After a 30 min incubation (37°C) of reaction mixtures containing the components described in Example 1, 75  $\mu$ M 2-methoxyestradiol was added, and aliquots were placed on 200-mesh carbon coated copper grids and stained with 0.5% (w/v) uranyl acetate. TEM magnifications from 23,100X to 115,400X

10 were used to visualize differences in tubulin morphology.

Example 3:

Figure 2 illustrates that 2-methoxyestradiol inhibits colchicine binding to tubulin. Reaction conditions were as described in the text, with each reaction mixture

15 containing 1.0  $\mu$ M tubulin, 5% (v/v) dimethyl sulfoxide, 5  $\mu$ M [<sup>3</sup>H]colchicine, and inhibitor at the indicated concentrations. Incubation was for 10 min at 37°C. Symbols as follows:  $\circ$ , 2-methoxyestradiol;  $\bullet$ , combretastatin A-4;  $\Delta$ , dihydrocombretastatin A-4. Combretastatin A-4 and

20 dihydrocombretastatin A-4 are compounds with anti-mitotic activity similar to colchicine.

Example 4:

Table 1 illustrates the inhibitory effects on tubulin polymerization in vitro exhibited by estradiol or

25 estradiol derivatives, plant anti-mitotic compounds such as colchicine, combretastatin A-4 or other plant compounds. The method is given in Example 1.

Example 5:

Table 2 lists estrogens, estradiol or estradiol

30 derivatives that inhibit colchicine binding to tubulin, by the method given in Example 3.

Table 1

	<u>Estrogenic Compound</u>	<u>IC<sub>50</sub> (μM ± S.D.)</u>
	2-Methoxyestradiol	1.9 ± 0.2
	Diethylstilbestrol	2.4 ± 0.4
5	2-Bromoestradiol ✓	4.5 ± 0.6
	2-Methoxyestrone	8.8 ± 1
	17-Ethynylestradiol	10.0 ± 2
	2-Fluoroestradiol	27.0 ± 6
	Estradiol	30.0 ± 6
10	Estrone	> 40
	2-Methoxy-17-ethynylestradiol ✓	> 40
	Estriol	> 40
	2-Methoxyestriol	> 40
	Estradiol-3-O-methyl ether ✓	> 40
15	2-Methoxyestradiol-3-O-methyl ether	> 40
	4-Methoxyestradiol	> 40
	4-Methoxyestradiol-3-O-methyl ether ✓	> 40
<hr/>		
	<u>Plant Products</u>	<u>IC<sub>50</sub> (μM ± S.D.)</u>
20	Colchicine	0.80 ± 0.07
	Podophyllotoxin	0.46 ± 0.02
	Combretastatin A-4	0.53 ± 0.05
	Dihydrocombretastatin A-4	0.63 ± 0.03
<hr/>		
25	IC <sub>50</sub> values are defined as the concentration of an estradiol derivative required to inhibit tubulin polymerization by 50%. IC <sub>50</sub> values were obtained in at least two independent experiments for non-inhibitory agents (IC <sub>50</sub> > 40 μM) and at least three independent experiments for inhibitory	
30	compounds. IC <sub>50</sub> values were obtained graphically, and average values are presented. S.D., standard deviation.	

2007142 021502

Table 2

	<u>Estrogenic Compound</u>	<u>Percent inhibition <math>\pm</math> S.D.</u>
	2-Methoxyestradiol	82 $\pm$ 2
	2-Methoxyestrone	57 $\pm$ 6
5	17-Ethynylestradiol	50 $\pm$ 7
	Estradiol	38 $\pm$ 4
	Diethylstilbestrol	30 $\pm$ 4

Reaction conditions were described in Example 3, with each  
10 reaction mixture containing 1.0  $\mu$ M tubulin, 5% (v/v)  
dimethyl sulfoxide, 2  $\mu$ M [ $^3$ H]colchicine, and 100  $\mu$ M  
inhibitor. Incubation was for 10 min at 37°C. Average  
values obtained in three independent experiments are  
presented in the table, except for 2-methoxyestrone, which  
15 was only examined twice. S.D., standard deviation.

What is claimed is: